



# Phylogenetic analysis of the Common Krait (*Bungarus caeruleus*) in Pakistan based on mitochondrial and nuclear protein coding genes

<sup>1</sup>Muhammad Rizwan Ashraf, <sup>1,\*</sup>Asif Nadeem, <sup>2</sup>Eric Nelson Smith, <sup>1</sup>Maryam Javed, <sup>2</sup>Utpal Smart, <sup>1</sup>Tahir Yaqub, <sup>1</sup>Abu Saeed Hashmi, and <sup>2</sup>Panupong Thammachoti

<sup>1</sup>Institute of Biochemistry and Biotechnology, University of Veterinary and Animal Sciences, Lahore, PAKISTAN <sup>2</sup>Amphibian and Reptile Diversity Research Center and Department of Biology, University of Texas at Arlington, Arlington, Texas 76019, USA

**Abstract.**—Pakistan has more than 40 species of venomous snakes. One of them, the Common Krait (*Bungarus caeruleus*), is responsible for most of the reported snake bites followed by Russel’s Viper, Saw-scaled Viper, and Black Cobra. Molecular studies not only help in correctly identifying organisms but also in finding the phylogenetic relationships and diversity among and between them. Morphological studies can be supplemented with confirmatory molecular data to make them more authentic and accurate. This study is the first to characterize the genetic diversity and phylogenetic relationships of Common Kraits from Pakistan, which will help in developing effective strategies for managing snake bites through effective antivenom development. Tail tip biopsies of 25 Common Kraits were collected from different cities in Pakistan. The whole DNA was extracted. Four mitochondrial (ND4, Cytochrome b, 12S rRNA, and 16S rRNA) and three nuclear protein coding (C-mos, RAG-1, and NT3) gene fragments were amplified using specific PCR primers. The amplified DNA was sequenced by Sanger di-deoxy sequencing. Forward and reverse sequences were cleaned and contiged using Sequencher 5.0 software. DNA data were aligned and concatenated using MEGA 6.0 and SequenceMatrix software, respectively. Partition Finder software was used for obtaining the best partitioning scheme and evolutionary models. Concatenated maximum likelihood and Bayesian phylogenetic trees were constructed using RaxML and MrBayes software. The same alignments were used to perform DNA polymorphism analysis using DnaSP 5.0 software. A percent identity matrix was created for all sequences using the online bioinformatics tool, MUSCLE. Homology was presented in tabular form, showing the similarity among different species of genus *Bungarus*. All *Bungarus* species were differentiated into four groups. Common Krait (*B. caeruleus*) from Pakistan showed close relationships with *B. sindanus* and *B. ceylonicus*, as one monophyletic group. The first clade included *B. candidus* (Indonesia, Thailand, Vietnam, and Laos), *B. multicinctus* (China, Taiwan, and Burma), and *B. niger* (Nepal). The second clade included *B. sindanus* and *B. caeruleus* (Pakistan), and *B. ceylonicus* (Sri Lanka). The third clade included *B. fasciatus* (Thailand and Indonesia), while the fourth clade included *B. bungroides* (China) and *B. flaviceps* (Malaysia and Indonesia). This study traces the diversity and phylogenetic relationships of the Pakistani elapid, Common Krait, showing the considerable inter- and intra-specific variations from different geographical regions of the world.

**Keywords.** Asia, Elapidae, PCR, polymorphism, Serpentes, venomous snakes

**Citation:** Ashraf MR, Nadeem A, Smith EN, Javed M, Smart U, Yaqub T, Hashmi AS, Thammachoti P. 2019. Phylogenetic analysis of the Common Krait (*Bungarus caeruleus*) in Pakistan based on mitochondrial and nuclear protein coding genes. *Amphibian & Reptile Conservation* 13(2) [General Section]: 203–211 (e205).

**Copyright:** © 2019 Ashraf et al. This is an open access article distributed under the terms of the Creative Commons Attribution License [Attribution 4.0 International (CC BY 4.0): <https://creativecommons.org/licenses/by/4.0/>], which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. The official and authorized publication credit sources, which will be duly enforced, are as follows: official journal title *Amphibian & Reptile Conservation*; official journal website: [amphibian-reptile-conservation.org](http://amphibian-reptile-conservation.org).

**Received:** 25 January 2019; **Accepted:** 24 September 2019; **Published:** 17 December 2019

## Introduction

Snakes are legless carnivorous reptiles of suborder Serpentes, and their lack of eyelids and external ears distinguish them from legless lizards (Reeder et al. 2015). Except Antarctica and some other large islands, like Ireland, Iceland, Greenland, the Hawaiian archipelago,

and New Zealand islands, snakes are found everywhere in the world (Roland 1994). South Asia is the region most affected with venomous snake bites. For example, the World Health Organization reports 35,000 to 50,000 deaths annually in India (Chippaux 1998; Pyron et al. 2013), and Pakistan reports 40,000 snake bites every year that result in 8,200 fatalities (Pyron et al. 2013).

**Correspondence.** \* [asifnadeem@uvas.edu.pk](mailto:asifnadeem@uvas.edu.pk)





**Fig. 1.** Sample collection sites in Pakistan for Common Krait (*Bungarus caeruleus*).

Venomous snakes in Southeast Asia belong to families Elapidae (cobras and kraits) and Viperidae (typical vipers and pit vipers). A study of hospital-admitted snakebite cases in Pakistan revealed less than 5% neurotoxic snakebites, and the rest were viper bites (Nisar et al. 2009).

Kraits, genus *Bungarus*, are identified by alternating black and white cross-bands across the body, and are found in all South Asian countries except the Philippines. Members of genus *Bungarus* are moderate to large sized elapids distributed in Pakistan and eastward through southern Asia to Indonesia (Smith 1943). Currently, 12 species of kraits are recognized (Yulin et al. 2018), including three in Pakistan. In Pakistan, Common Kraits (*Bungarus caeruleus*) are reported throughout Punjab, Khyber Pakhtoonkhwa (KPK), Azad Kashmir, Sindh, and Southern Balochistan. Common in the Indus valley, this is the only species of kraits found in Rawalpindi and Islamabad (Khan 2002a; Oh et al. 2019). Sindh Krait (*B. sindanus*) is prevalent in Tharparkar, Bahawalnagar, and Bahawalpur. Northern Punjab Krait (*B. razai*) is reported from Mianwali (Khan 2002b). Determining the relationships among the members of Elapidae can help in understanding their distribution and diversity. Many studies have focused on the evolutionary relationships of kraits.

Before the widespread use of DNA sequencing, systematics and taxonomy were used to infer phylogenies among species in order to explain their relationships. Now many fields in biology are using phylogenies for a wide variety of purposes, such as examining paralogous relationships, population histories, dynamics of pathogens with respect to their evolution and epidemiology (Zhou et al. 2018; Blanquart 2019), the ontogeny of body cells during development, and the differentiation of tumors (Kester and van Oudenaarden 2018). Variations in nucleotide sequences can construct phylogenies for

inferring relationships among the compared sequences. The topology of phylogeny gives some estimates about the mutation rates, time-scales of evolutionary events, and prehistoric movement among different geographical regions (Soroka and Burzyński 2018). Phylogenetics shows relationships among organisms and genes (Friberg et al. 2019), and can give a clearer picture of the biodiversity, biogeography, and evolution of many characters in related groups (Pilfold et al. 2019; Grismer and Davis 2018; Silva et al. 2019).

The use of mitochondrial DNA data for studying animal evolution has become a powerful tool in the last decade. Molecular biology has helped in these mitochondrial DNA studies to give insights into population structure, gene flow, hybridization, biogeography, and phylogenetics (Chandrasekaran et al. 2019; Soroka and Burzyński 2018). Evolutionary studies give comparisons of mitochondrial genome organization and function while molecular studies help to improve these evolutionary studies (Ng et al. 2019). Nuclear encoded genes seem to be a strong source of phylogenetic information. They can be more useful for showing the divergence of those genes whose multiple substitutions may obscure clear phylogenetic signals.

This is the first study from Pakistan focusing on genetic characterization, biodiversity, and molecular phylogenetics of the Common Krait (*Bungarus caeruleus*).

## Materials and Methods

A collection of 25 Common Kraits (*Bungarus caeruleus*) was obtained from reptile breeders in different cities in Pakistan (see Fig. 1, Table 1). Scalation patterns and

**Table 1.** Common Krait (*Bungarus caeruleus*) samples used in this study, and their location information.

Sample ID	Locality	Latitude	Longitude
BC-1	Jallo Park, Lahore, Punjab, Pakistan	31°34'17.29"N	74°28'36.78"E
BC-2	Balochanwali, Bahawalpur, Punjab, Pakistan	29°28'56.90"N	71°59'41.86"E
BC-3	Qila Ram Qaur, Hafiz Abad, Punjab, Pakistan	32°4'57.12"N	73°40'49.02"E
BC-4	Yazman Housing Society, Yazman, Punjab, Pakistan	29°7'3.00"N	71°45'6.73"E
BC-5	Changa Manga Forest, Kasur, Punjab, Pakistan	31°4'54.19"N	73°59'53.49"E
BC-6	Lal Suhanra National Park, Bahawalpur, Punjab, Pakistan	29°19'1.36"N	71°54'16.43"E
BC-7	Rahim Yar Khan Zoo, Rahim Yar Khan, Punjab, Pakistan	28°24'14.30"N	70°15'32.63"E
BC-8	Chak Risalwala, Faisalabad, Punjab, Pakistan	31°22'4.90"N	73°1'24.80"E
BC-9	Qila Ram Qaur, Hafizabad, Punjab, Pakistan	32°4'57.12"N	73°40'49.02"E
BC-10	New City Housing Society, Jaranwala, Punjab, Pakistan	31°19'16.60"N	73°23'21.44"E
BC-11	Chak 126 GB Pind Janjua, Jaranwala, Punjab, Pakistan	31°21'38.48"N	73°25'28.74"E
BC-12	Rahim Yar Khan Zoo, Rahim Yar Khan, Punjab, Pakistan	28°24'14.30"N	70°15'32.63"E
BC-13	Yazman Housing Scheme, Yazman, Punjab, Pakistan	29°6'54.39"N	71°45'17.40"E
BC-14	Ayub National Park, Jehlam Road, Punjab, Pakistan	33°34'19.00"N	73°4'59.00"E
BC-15	Tibbi Balochan, Sadiqabad, Punjab, Pakistan	28°16'35.01"N	70°8'6.58"E
BC-16	Maraghzar Colony, Lahore, Punjab, Pakistan	31°30'8.71"N	74°14'55.48"E
BC-17	Lahore Zoo, Punjab, Pakistan	31°33'23.78"N	74°19'33.73"E
BC-18	Lahore Zoo, Punjab, Pakistan	31°33'23.78"N	74°19'33.73"E
BC-19	Raza Garden Phase 1, Sargodha, Punjab, Pakistan	32°2'51.23"N	72°37'31.68"E
BC-20	Pir wala, Jhang, Punjab, Pakistan	31°1'42.61"N	72°16'45.51"E
BC-21	Noor Garden, Okara, Punjab, Pakistan	30°48'48.38"N	73°28'38.33"E
BC-22	Chenab Park, Multan, Punjab, Pakistan	30°4'29.90"N	71°18'51.93"E
BC-23	Kalarwala, Chiniot, Punjab, Pakistan	31°28'26.21"N	72°33'56.11"E
BC-24	Qadir Abad Tiba, Sadiqabad, Punjab, Pakistan	28°16'54.33"N	70°7'45.48"E
BC-25	Chenab Park, Gujranwala, Punjab, Pakistan	30°4'29.90"N	71°18'51.93"E


tail tip biopsies were obtained from each specimen. After DNA extraction (Sambrook and Russel 2001), the Polymerase Chain Reaction (PCR) primers of representative mitochondrial genes (ND4, Cytochrome b, 12S rRNA, and 16S rRNA) and nuclear genes

(C-mos, RAG1, and NT3) from previous studies were used for the amplification of selected regions through PCR (see Table 2). After amplification, amplicons were sequenced bi-directionally by Big Dye™ Terminator on an ABI 3130XL Genetic analyzer. Forward and

**Table 2.** Mitochondrial and nuclear protein coding gene primers for Common Krait (*Bungarus caeruleus*).

Sr. No	Gene Name	Primer Sequence	Source
1	Cyt.b	5'-TGACTTGAARAACCAAYCGTTG-3'	Palumbi 1996
		5'-TGAGAAGTTTTTCYGGGTCRRT-3'	Parkinson et al. 2002
2	16S rRNA	5'-CGCCTGTTTAYCAAAAACAT-3	Vences et al. 2005
		5'-CCGGTCTGAACTCAGATCACGT-3'	Vences et al. 2005
3	12S rRNA	5'GTACACTTACCTTGTTACGACTT 3'	Knight and Mindell 1993
		5'AACTGGGATTAGATACCCCACTAT3'	Knight and Mindell 1993
4	ND4	5'-CATTACTTTTACTTGGATTGACCA-3'	Arevalo 1994
		5'-CACCTATGACTACCAAAAGCTCATGTAAGC-3'	Arevalo 1994
5	RAG-1	5'AGCTGCAGYCARTAYCAYAARATGTA3'	Chiari et al. 2004
		5'AACTCAGCTGCATTKCCAATRTCA3'	Chiari et al. 2004
6	NT3	5'ATATTTCTGGCTTTTCTCTGTGGC3'	Townsend et al. 2008
		5'GCGTTTCATAAAAATATTGTTTGACC3'	Townsend et al. 2008
7	C-mos	5'CATGGACTGGGATCAGTTATG 3'	Lawson et al. 2005
		5'CCTTGGGTGTGATTTTCTCACCT 3'	Lawson et al. 2005





**Fig. 2.** Mitochondrial and nuclear genes (ND4, Cyt. b, COI, 12S rRNA, 16S rRNA, C-mos, RAG-1, NT3, and BDNF) based Maximum Likelihood phylogeny for Common Krait (*Bungarus caeruleus*).

reverse sequences were assembled through Sequencher 5.0 software. The resulting contigs (sequences) were given specific identities. These contigs were then aligned with other reported sequences obtained from the NCBI database through MEGA (v 6.0, Tamura et al. 2013) using the ClustalW tool for further data analyses. The nucleotide data for each gene were concatenated using SequenceMatrix (v1.7.8, Vaidya et al. 2011) software. The concatenated data were partitioned through PartitionFinder (v1.1.1, Lanfear et al. 2012) to give the best partition scheme and evolutionary models for phylogenetic analyses.

Two types of phylogenetic analyses, i.e., Maximum Likelihood (ML) and Bayesian inference (BI), were performed through RaxML (v8.0, Stamatakis 2014) and MrBayes (v3.2, Ronquist and Huelsenbeck 2012) software. The resulting phylogenetic trees were visualized and saved using Figtree (v1.4.3, <http://tree.bio.ed.ac.uk/software/figtree/>) software. DnaSP (v5.0, Librado and Rozas 2009) was used for analyzing polymorphic sites and DNA polymorphism, to determine the variation and genetic biodiversity in Common Krait (*Bungarus caeruleus*) in relation to other species of the genus *Bungarus*. Percent identity matrices were also constructed by comparing different species of every snake genus using online tool MUSCLE (available from the European Molecular Biology Laboratory, <https://www.ebi.ac.uk/Tools/msa/muscle/>).


## Results

DnaSP software was used for analyzing the polymorphism of the mitochondrial and nuclear genes as shown in Table 3. The ribosomal RNA coding genes showed the least

variation, with lower numbers of variable sites, mutations, and parsimony informative sites. The polymorphism data show the variations in different mitochondrial and nuclear genes with their conservation among Common Krait and other species of genus *Bungarus*. In addition, no significant variations were found among Common Kraits from different cities in Pakistan.

The online MUSCLE tool was used to find relationships among *Bungarus* species on the basis of homology in the mitochondrial and nuclear genes (Table 4). This table shows the conservation patterns in mitochondrial and nuclear protein coding genes of various *Bungarus* species.

Phylogenetic analysis of Common Krait (*Bungarus caeruleus*) from Pakistan was conducted using mitochondrial and nuclear protein coding genes. In this study, Black Cobra (*Naja naja*) from Thatta Sindh was used as the outgroup for constructing maximum likelihood and Bayesian phylogenies. The best partition scheme and evolutionary models were used to infer the phylogenetic relationships of Common Krait in Pakistan with other members of genus *Bungarus* around the world. Concatenated Maximum likelihood and Bayesian Inference results gave very similar phylogenies (Figs. 2–3). All *Bungarus* species were divided into four main clades. The first clade included *B. candidus* (Indonesia, Thailand, Vietnam, and Laos), *B. multicinctus* (China, Taiwan, and Burma), and *B. niger* (Nepal). The second clade included *B. sindanus* and *B. caeruleus* (Pakistan), and *B. ceylonicus* (Sri Lanka). The third clade included *B. fasciatus* (Thailand and Indonesia), while the fourth clade included *B. bungaroides* (China) and *B. flaviceps* (Malaysia and Indonesia). The first and second clades showed a sister clade relationship with strong support (ML BS = 100, BI PP = 1). *Bungarus candidus* and



**Fig. 3.** Mitochondrial and nuclear genes (ND4, Cyt b, COI, 12S rRNA, 16S rRNA, C-mos, RAG-1, and NT3) Bayesian phylogeny for Common Krait (*Bungarus caeruleus*).

*B. multicinctus* probably diverged as separate species only recently. *Bungarus candidus*, *B. multicinctus*, and *B. niger* showed highly supported sister relationships with a complex pattern of divergence (BI PP = 1.0 ML BS = 90). In the second clade *B. sindanus* and *B. caeruleus* have been reported from Pakistan, thus they are sympatric species, while *B. ceylonicus* (from Sri Lanka) also showed a significant difference with strong support through Maximum likelihood and Bayesian inference phylogenies (PP = 1.0 and BS = 100). Pyron et al. (2012) also revealed the same relationships between *B. caeruleus*, *B. sindanus*, and *B. ceylonicus*.

In addition to the molecular data obtained, examination of the 25 *B. caeruleus* specimens showed varying numbers of ventral scales (207–218), 15 rows of mid-body scales, and average numbers of subcaudals of 41–47.

## Discussion

Elapids comprise 300 of the 2,500 known species of snakes (Leviton et al. 2018). The Southern Asian

elapids include cobras (*Naja* and *Ophiophagus*), kraits (*Bungarus*), long-glanded snakes (*Maticora*), and Asian coral snakes (*Calliophis*) [Sanz et al. 2019]. The uncertain phylogenetics of elapids has been a major factor for the varying numbers of identified species of elapids in the past (Mirtschin et al. 2017).

This study aimed to characterize the genetic biodiversity and phylogenetic relationships of Common Krait (*Bungarus caeruleus*) as there is a great deal of unpublished data on this species. Here, mitochondrial and nuclear protein coding genes were used to construct the phylogeny of Common Krait from Pakistan along with some morphological characterization. One variable character is the number of ventral scales that ranges from 207–218 among the 25 specimens in this study. Khan (1985) wrote a note on the taxonomic status of Common Krait and Sindh Krait (*B. sindanus*), and by comparing 46 specimens, Khan noted an almost similar range of 207–218 ventral scales. The *B. caeruleus* in this study had 15 rows of mid-body scales which is the same as described by Khan (1985). *Bungarus caeruleus* and *B.*

**Table 3.** Polymorphism in mitochondrial and nuclear protein coding genes of Common Krait (*Bungarus caeruleus*).

Parameters	ND4	Cytochrome b	12S rRNA	16S rRNA	C-mos	RAG-1	NT3
Total number of sites	619	702	650	520	586	802	425
Variable number of sites	302	325	270	191	419	665	325
Number of mutations	302	202	63	47	12	20	32
Singleton variable sites	43	51	20	26	09	19	03
Parsimony informative sites	196	151	43	21	03	01	28
Segregating sites	159	151	00	00	12	19	31
Synonymous changes	176	156	00	00	06	03	23
Number of haplotypes	17	16	08	10	04	03	05
Haplotype diversity	0.866	0.615	0.686	0.521	0.333	0.145	0.754
Nucleotide diversity	0.08243	0.09528	0.03617	0.03451	0.00288	0.00226	0.03389

**Table 4.** Percent homology of mitochondrial and nuclear genes for Common Krait (*Bungarus caeruleus*) from Pakistan among various *Bungarus* species and countries, sorted by decreasing Cytb homology. The asterisk (\*) indicates the sequence from this study, all others are based on GenBank sequences.

Species	Country	Homology percentages of nine representative genes								
		Cytb	ND4	12S	16S	COI	C-mos	RAG-1	NT3	BDNF
<i>B. caeruleus</i> *	Pakistan	100	100	100	100	100				
<i>B. caeruleus</i>	Pakistan	99.84	99.51	100	100	99.99				
<i>B. ceylonicus</i>	NA	89.21	87.46	93.68	96.62		100	99.4		
<i>B. candidus</i>	Indonesia	89.19	84.53	90.57	95.08					
<i>B. niger</i>	Nepal	86.31	84.36							
<i>B. sindanus</i>	Pakistan	86.31	85.5							
<i>B. candidus</i>	Thailand	85.35	83.88							
<i>B. multicinctus</i>	Burma	85.19	84.34			85.65				
<i>B. multicinctus</i>	China	85.19	85.18							
<i>B. candidus</i>	Vietnam	85.19	85.67							
<i>B. multicinctus</i>	Taiwan	85.02	85.5	91.73	90.09	86.11				
<i>B. fasciatus</i>	Thailand	83.74	83.55	90.79	91.91					
<i>B. fasciatus</i>	Indonesia	83.57	83.39	91.38	95.5	85.03	99.38	99.62	96.63	99.76

*sindanus* had 15 and 17 rows of mid-body scales, with the central larger row being hexagonal and white in color. Boulenger (1897) also observed 15 rows of mid-body scales in Indo-Pakistan Common Krait while 17 mid-body scale rows were reported in *B. sindanus* from Indus Basin. The average number of sub-caudals observed here is 41–47 which is within the range observed by Khan (1985): 40–54 in males and 30–54 in females. Boulenger (1897) reported small eyes with round pupil which is similar to those observed in this study.

There are also reports about the distribution of *B. caeruleus* in Indo-Pakistan subcontinent. Eastward it is found in Assam and Bengal (Jamal et al. 2018; Ganesh and Vogel 2018); westward to the Pakistan-Iran Border; Shockley (1949) and Kral (1969) reported it in Afghanistan; Smith (1943) reported it southward in Peninsular India and the Andaman Islands; and de Silva (1981) also reported it in Sri Lanka. This study is one attempt to infer the phylogenetics of *B. caeruleus* in Pakistan, but suggests more studies from the above-mentioned parts of the *B. caeruleus* distribution are needed, as there are almost no studies from other parts of the Indo-Pakistan subcontinent on the phylogenetics of the Common Krait *B. caeruleus*.

Determining the relationships among the members of Elapidae can help in understanding the distribution and diversity of elapids. Many studies have focused on evolutionary relationships of elapids, and this study examined the molecular phylogenetics of *B. caeruleus* from Pakistan. The second clade (Figs. 2–3) includes *B. sindanus* and *B. caeruleus* (Pakistan), and *B. ceylonicus* (Sri Lanka). *Bungarus sindanus* and *B. caeruleus* are sympatric species, while *B. ceylonicus* also showed a significant difference with strong support through the Maximum likelihood and Bayesian inference phylogenies

(PP = 1.0 and BS = 100). Pyron et al. (2012) also revealed the same relationships between *B. caeruleus*, *B. sindanus*, and *B. ceylonicus*. They presented a large-scale phylogeny of squamate reptiles for future comparative studies, and a revised classification of squamates at the family and subfamily levels so that taxonomy might be brought in a line with data from the new phylogenetic studies. Their phylogeny shows the same relationship (ML BS = 100, BI = 1) between *B. caeruleus*, *B. sindanus*, and *B. ceylonicus* as is shown in this study through Maximum Likelihood and Bayesian phylogenies.

## Conclusions

Most of the currently recognized krait species (genus *Bungarus*) are poorly understood. This study characterized the genetic biodiversity and phylogenetic relationships of Common Krait (*Bungarus caeruleus*) from Pakistan showing inter- and intra-specific variations among different geographical regions of the world. More diverse sampling and a larger number of samples with more genomic data could help to further resolve the taxonomic status of the *Bungarus* species in Pakistan. This study also provides guidance for the correct identification of these snakes with authentication using molecular biology tools which will be helpful in the development of effective and region-specific antivenoms for such venomous snakes.

## Literature Cited

Arevalo E, Davis SK, Sites JW Jr. 1994. Mitochondrial DNA sequence divergence and phylogenetic relationships among eight chromosome races of the *Sceloporus grammicus* complex (Phrynosomatidae)



- in Central Mexico. *Systematic Biology* 43(3): 387–418.
- Blanquart F. 2019. Evolutionary epidemiology models to predict the dynamics of antibiotic resistance. *Evolutionary Applications* 12(3): 365–383.
- Boulenger GA. 1897. A new krait from Sind (*Bungarus sindanus*). *Journal of the Bombay Natural History Society* 11: 73–74.
- Chandrasekaran K, Anjaneyulu M, Choi J, Kumar P, Salimian M, Ho CY, Russell JW. 2019. Role of mitochondria in diabetic peripheral neuropathy: Influencing the NAD<sup>+</sup>-dependent SIRT1-PGC-1 $\alpha$ -TFAM pathway. *International Review of Neurobiology* 145: 177–209.
- Chiari Y, Vences M, Vieites DR, Rabemananjara F, Bora P, Ravoahangimalala OR, Meyer A. 2004. New evidence for parallel evolution of color patterns in Malagasy poison frogs (*Mantella*). *Molecular Ecology* 13: 3,763–3,774.
- Chippaux JP. 1998. Snake-bites: appraisal of the global situation. *Bulletin of the World Health Organization* 76: 515–524.
- da Silva WO, da Costa MJR, Pieczarka JC, Rissino J, Pereira JC, Ferguson-Smith MA, Nagamachi CY. 2019. Identification of two independent X-autosome translocations in closely related mammalian (*Proechimys*) species. *Scientific Reports* 9(1): 4,047.
- De Silva A. 1981. Snakebites in Anuradhapura district. *The Snake* 13: 117–130.
- Friberg M, Schwann C, Guimarães PR, Raguso RA, Thompson JN. 2019. Extreme diversification of floral volatiles within and among species of *Lithophragma* (Saxifragaceae). *Proceedings of the National Academy of Sciences of the United States of America* 116(10): 4,406–4,415.
- Ganesh SR, Vogel G. 2018. Taxonomic reassessment of the Common Indian Wolf Snakes *Lycodon aulicus* (Linnaeus, 1758) complex (Squamata: Serpentes: Colubridae). *Bonn Zoological Bulletin* 67(1): 25–36.
- Grismer LL, Davis HR. 2018. Phylogeny and biogeography of Bent-toed Geckos (*Cyrtodactylus* Gray) of the Sundaic swamp clade. *Zootaxa* 4472(2): 365–374.
- Hoffstetter R, Gasc JP. 1969. Vertebrae and ribs of modern reptiles. Pp. 201–310 In: *Biology of the Reptilia. Volume 1: Morphology*. Editors, Gans C, Bellairs A, Parsons TS. Academic Press, London, United Kingdom. 373 p.
- Jamal Q, Idrees M, Ullah S, Adnan M, Zaidi F, Zaman Q, Rasheed SB. 2018. Diversity and altitudinal distribution of Squamata in two distinct ecological zones of Dir, a Himalayan sub-zone of northern Pakistan. *Pakistan Journal of Zoology* 50(5): 1,835–1,839.
- Kester L, van Oudenaarden A. 2018. Single-cell transcriptomics meets lineage tracing. *Cell Stem Cell* 23(2): 166–179.
- Khan MS. 1985. Taxonomic notes on *Bungarus caeruleus* (Schneider) and *Bungarus sindanus* Boulenger. *The Snake* 17: 71–78.
- Khan MS. 2002a. *Die Schlangen Pakistans*. Edition Chimaira, Frankfurt am Main, Germany. 265 p.
- Khan MS. 2002b. *A Guide to the Snakes of Pakistan*. Edition Chimaira, Frankfurt am Main, Germany. 265 p.
- Knight A, Mindell DP. 1993. Substitution bias, weighting of DNA sequence evolution, and the phylogenetic position of Fea's Viper. *Systematic Biology* 42: 18–31.
- Kral B. 1969. Notes on the herpetofauna of certain provinces of Afghanistan. *Zoologische Listy* 18: 55–66.
- Lanfear R, Calcott B, Ho SY, Guindon S. 2012. PartitionFinder: combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and Evolution* 29: 1,695–1,701.
- Lawson R, Slowinski JB, Crother BI, Burbrink FT. 2005. Phylogeny of the Colubroidea (Serpentes): new evidence from mitochondrial and nuclear genes. *Molecular Phylogenetics and Evolution* 37: 581–601.
- Leviton AE, Siler CD, Weinell JL, Brown RM. 2018. Synopsis of the snakes of the Philippines. *Proceedings of the California Academy of Sciences* 64(14): 399–568.
- Librado P, Rozas J. 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25(11): 1,451–1,452.
- Miller MA, Pfeiffer W, Schwartz T. 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. Pp. 1–8 In: *2010 Gateway Computing Environments Workshop*. Institute of Electrical and Electronics Engineers, Piscataway, New Jersey, USA.
- Mirtschin P, Rasmussen A, Weinstein S. 2017. *Australia's Dangerous Snakes: Identification, Biology and Envenoming*. CSIRO Publishing, Melbourne, Australia. 432 p.
- Ng YS, DiMauro S, Turnbull DM. 2019. Mitochondrial medicine: a historical point of view. Pp. 1–18 In: *Diagnosis and Management of Mitochondrial Disorders*. Editors, Mancuso M, Klopstock T. Springer, Cham, Switzerland. 382 p.
- Nisar A, Rizvi F, Afzal M, Shafi MS. 2009. Presentation and complications of snakebite in a tertiary care hospital. *Journal of College of Physicians and Surgeons Pakistan* 19: 304–307.
- Oh AMF, Tan CH, Tan KY, Quraishi NH, Tan NH. 2019. Venom proteome of *Bungarus sindanus* (Sind krait) from Pakistan and in vivo cross-neutralization of toxicity using an Indian polyvalent antivenom. *Journal of Proteomics* 193: 243–254.
- Palumbi SR. 1996. Nucleic acids II: the polymerase chain reaction. Pp. 205–247 In: *Molecular Systematics*. 2<sup>nd</sup> Edition. Editors, Hillis DM, Moritz C, Mable BK. Sinauer Associates, Sunderland, Massachusetts, USA. 665 p.
- Parkinson CL, Campbell JA, Chippendale PT. 2002. Multigene phylogenetic analysis of pitvipers with comments on the biogeographical history of the

- group. Pp. 93–110 In: *Biology of the Vipers*. Editors, Schuett GW, Hoggren M, Douglas ME, Greene HW. Eagle Mountain Publishing, Utah, USA. 580 p.
- Pilfold NW, Letoluai A, Ruppert K, Glikman JA, Stacy-Dawes J, O'Connor D, Owen M. 2019. Confirmation of black leopard (*Panthera pardus pardus*) living in Laikipia County, Kenya. *African Journal of Ecology* 57(2): 270–273.
- Pyron RA, Burbrink FT. 2012. Extinction, ecological opportunity, and the origins of global snake diversity. *Evolution* 66: 163–178.
- Pyron RA, Burbrink FT, Wiens JJ. 2013. A phylogeny and revised classification of Squamata, including 4,161 species of lizards and snakes. *BMC Evolutionary Biology* 13: 93.
- Reeder TW, Townsend TM, Mulcahy DG, Noonan BP, Wood Jr PL, Sites Jr JW, Wiens JJ. 2015. Integrated analyses resolve conflicts over squamate reptile phylogeny and reveal unexpected placements for fossil taxa. *PLOS One* 10: e0118199.
- Roland B. 1994. *Snakes: A Natural History*. Sterling Publishing, New York, New York, USA. 220 p.
- Ronquist F, Huelsenbeck JP. 2012. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1,572–1,574.
- Sambrook J, Russel DW. 2001. Rapid isolation of yeast DNA. Pp. 631–632 In: *Molecular Cloning, a Laboratory Manual*. Editors, Sambrook J, Russel, DW. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, USA. 721 p.
- Sanz L, Quesada-Bernat S, Ramos T, Casais-e-Silva LL, Corrêa-Netto C, Silva-Haad JJ, Calvete JJ. 2019. New insights into the phylogeographic distribution of the 3FTx/PLA2 venom dichotomy across genus *Micrurus* in South America. *Journal of Proteomics* 200: 90–101.
- Shockley CH. 1949. Herpetological notes for Ras Jiunri, Baluchistan. *Herpetologica* 5: 121–123.
- Smith MA. 1943. *The Fauna of British India including Ceylon and Burma*. Taylor and Francis, London, United Kingdom. 185 p.
- Soroka M, Burzyński A. 2018. Doubly uniparental inheritance and highly divergent mitochondrial genomes of the freshwater mussel *Unio tumidus* (Bivalvia: Unionidae). *Hydrobiologia* 810(1): 239–254.
- Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1,312–1,313.
- Tamura K, Stecher G, Peterson D, Filipowski A, Kumar S. 2013. MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. *Molecular Biology and Evolution* 12: 2,725–2,729.
- Townsend TM, Alegre RE, Kelly ST, Wiens JJ, Reeder TW. 2008. Rapid development of multiple nuclear loci for phylogenetic analysis using genomic resources: an example from squamate reptiles. *Molecular Phylogenetics and Evolution* 47: 129–142.
- Vaidya G, Lohman DJ, Meier R. 2011. SequenceMatrix: concatenation software for the fast assembly of multi-gene datasets with character set and codon information. *Cladistics* 27: 171–180.
- Vences M, Thomas M, Bonett RM, Vieites DR. 2005. Deciphering amphibian diversity through DNA barcoding: chances and challenges. *Philosophical Transactions of the Royal Society of London B. Biological Sciences* 360(1462): 1,859–1,868.
- Yulin X, Ping W, Guanghui Z. 2018. Molecular phylogeny found the distribution of *Bungarus candidus* in China (Squamata: Elapidae). *Zoological Systematics* 43(1): 109–117.
- Zhou P, Fan H, Lan T, Yang XL, Shi WF, Zhang W, Zheng XS. 2018. Fatal swine acute diarrhoea syndrome caused by an HKU2-related coronavirus of bat origin. *Nature* 556(7700): 255.




**Muhammad Rizwan Ashraf** is a scholar at the University of Veterinary and Animal Sciences, Lahore, Pakistan. Muhammad has an M.Sc. in Biochemistry and Ph.D. in Molecular Biology and Biotechnology. His research interest is reptilian (especially snake) molecular phylogeny and genetic biodiversity. Muhammad visited University of Texas at Arlington, Texas, USA as a research scholar under the International Research Support Initiative Program, funded by the Government of Pakistan, and his Ph.D. work involved four of the most venomous snakes of Pakistan: Black Cobra, Common Krait, Saw-scaled Viper, and Russell's Viper. He is seeking further challenges to expand his skills within a progressive and fast-growing environment that uses state-of-art technologies while enhancing his proven abilities and dedication to team motivation.



**Asif Nadeem** is an Associate Professor at the University of Veterinary and Animal Sciences, Lahore, Pakistan. Asif had the privilege of receiving research training from the University of Wisconsin–Madison, Wisconsin, USA, and he has maintained a vigorous research program in the genetics of animals of agricultural importance. His research has generated many publications in peer-reviewed journals, and he has presented his research work at various national and international forums. Dr. Nadeem's efforts were recognized by his receipt of the Research Productivity Award from the Pakistan Commission of Science and Technology. He is an internationally known researcher in the field of animal genetics and genomics, and has served as an editor and reviewer for many different scientific journals.





**Eric Nelson Smith** is a Professor at University of Texas at Arlington, Texas, USA. Eric's current research interest focuses on the *Exploration and Speciation in the Volcanoes of the Indonesian Ring of Fire: a Large-Scale Inventory of the Herpetofauna of the Highlands of Sumatra and Java*. As a benefit to the scientific community, this project is producing modern specimen repositories in the two participating countries and web-based resources for identification and conservation, as well as for genetic and biodiversity work. Many of these efforts have been directed toward the systematics and taxonomy of venomous snakes. Eric has participated in the *Threatened Amphibians of the World* project (<https://portals.iucn.org/library/node/9186>) and the International Union for the Conservation of Nature (IUCN) Red List Assessment of Reptile Species, particularly with species of Asian coral snakes.



**Maryam Javed** is an Assistant Professor at the University of Veterinary and Animal Sciences, Lahore, Pakistan. She has received many academic and scientific awards, including a gold medal in D.V.M., Star Laureate Award, Nestle Award, Tufail Muhammad Award, Best Student in Academics Award from the Chancellor, Governor of Punjab, and the all Pakistan Quid Talent Award. Maryam's current research interest focuses on the identification of genes of economic importance in dairy animals.



**Utpal Smart** hails from Pondicherry, a sleepy coastal town (of *Life of Pi* fame) in southern India. Utpal's formal training in biology began with an M.Sc. in Ecology from Pondicherry University, India, in 2008, followed by a Ph.D. in Quantitative Biology from the University of Texas at Arlington, USA, in 2016. His doctoral training primarily involved using computational methods to investigate questions in molecular ecology using a combination of macro- and micro-evolutionary approaches. As a postdoctoral research associate at University of North Texas Center for Human Identification (UNTCHI), Utpal is helping to create the Mitochondrial Mixture Database and Interpretation Tool (MMDIT)—a bioinformatic pipeline for deconvoluting mitochondrial DNA mixtures and using computational phylogenetic and population genetic methods on human microbiome data to leverage them as forensic tools.



**Tahir Yaqub** is a Senior Member of the University of Veterinary and Animal Sciences (UVAS), Lahore, Pakistan, and has over 25 years of scientific experience in controlling infectious diseases of livestock, including those caused by influenza viruses. Tahir did a postdoc at the Institute of Public Health, United Kingdom, and his research interests include investigating the biological health risks of various agents in public and animal health. His laboratory has postgraduate students with expertise in molecular biology and serves as a hub for training. Currently, Tahir is investigating the prevalence and control of various diseases of public health importance.



**Abu Saeed Hashmi, Ph.D.**, is a well-known academician, researcher, and mentor in the field of Biochemistry. Abu's major area of research is in the bioconversion of agricultural/industrial waste to value added products. Abu has supervised many postgraduate students and has contributed significantly to this field. His work on bioconversion, Alfa toxins, bio-wastes, and production of biomass has been published and widely cited in many prominent research journals.



**Panupong Thammachoti** graduated from the University of Texas at Arlington, USA, in the Quantitative Biology Program (Ecology and Systematics). Panupong's work used multiple approaches including morphology, molecular phylogeny, and ecology, for solving taxonomic problems. As a lecturer at Chulalongkorn University, Thailand, he is interested in several research topics focusing on the diversity of amphibians and reptiles. Panupong has several scientific recognitions, such as a scholarship from the Human Resource Development in Science Project (Science Achievement Scholarship of Thailand, SAST); International Training Course-New Trends and Methodology in Animal Ecology and Conservation Biology; International Society of Zoological Sciences, Beijing, China; The Professor Dr. Tab Nilanidhi Foundation Award for outstanding academics, Faculty of Science, Chulalongkorn University; and the best oral presentation award, JSPS CORE-TO-CORE PROGRAM at the 5th International Symposium on Asian Vertebrate Species Diversity, Thailand.